

## HOST PLANT RANGE AND MORPHOMETRICS DESCRIPTIONS OF AN EMERGING INSECT PEST OF CASHEW, *Plocaederus ferrugineus* L. IN NIGERIA: A PRELIMINARY REPORT<sup>1</sup>

E. U. ASOGWA<sup>2</sup>, J. C. ANIKWE<sup>2</sup>, T. C. N. NDUBUAKU<sup>2</sup>, F. A. OKELANA<sup>2</sup> AND L. A. HAMMED<sup>2</sup>

<sup>2</sup> Cashew Research Programme, Cocoa Research Institute of Nigeria, Ibadan, Nigeria

Accepted for publication April 15, 2009

### ABSTRACT

Asogwa E. U., Anikwe J. C., Ndubuaku T. C. N., Okelana F. A. and Hamed L. A. 2009. *Host plant range and morphometrics descriptions of an emerging insect pest of cashew, plocaederus ferrugineus l. in nigeria: a preliminary report.* J. Innov. Dev. Strategy. 3(2):21-26

A 2.5 hectare cashew germplasm plot (S/GP) within Cocoa Research Institute of Nigeria, Ibadan was assessed between 2006 to 2007 to identify the host plant range and to describe the morphology of the various life stages of an emerging insect pest of cashew, *Plocaederus ferrugineus* in Nigeria. The cashew trunk and root borer, *Plocaederus ferrugineus* is fast becoming the most dreaded insect pest of cashew plantations in Nigeria as its infestation results in "sudden death" of mature cashew trees within few weeks. The rate of infestation of the borer on cashew and all tree stands/shrubs within S/GP and adjacent plots were recorded. Measurements of the morphometric characteristics of the various life stages of the borer collected from infested cut down trunks and excavated roots were carried out with a pair of caliper. Fewer stands of cashew were affected by the trunk and root borer resulting in an infestation rate of 36.3%. Only *Spondias mombin* was found to be riddled with borers out of all the other 24 species of trees and shrubs found within S/GP. 89.6% of the *S. mombin* assessed came down with different levels of attack by the borer, which differed significantly ( $P < 0.05$ ) from 36.3% infestation rate on cashew. The larva (grub) has a mean body length of 2.72mm, while an average pupal length of 3.66mm was recorded. The adult *P. ferrugineus* is a medium sized dark grey beetle, 2.91cm long and 0.92cm wide at the base of the abdomen. They have longer antennae (3.58cm), which is significantly longer ( $P < 0.05$ ) than their body (2.91cm).

**Key words:** Infestation, larvae, pupae, borer, adults, damages, germplasm.

### INTRODUCTION

The commercial cashew plantations started in Nigeria in the early 1950s with the establishment of first commercial plantations at Oghe, Oji and Mbala by the defunct Eastern Nigeria Development Corporation (ENDC) and Iwo, Eruwa and Upper Ogun by the defunct Western Nigeria Development Corporation (WNDC). From these locations, the planting of the crop spread to other parts of Nigeria particularly the Central and Northern States of Nigeria (Ohler, 1967; 1979; Venkataramah, 1976; Togun, 1977). The introduction of Brazilian cashew biotype with improved and desirable nut and kernel quality characteristics by CRIN has further increased the crops spread and popularity in Nigeria (Hamed *et al.*, 2007).

The cashew apple is sweet and when fully ripe, the juice is rich in vitamin C and sugars. It contains 7 - 9 % of reducing sugars, 11-12% of soluble solids and about 0.5% of tannic acid (Opeke, 1987). In most cashew producing nations, it is eaten raw. It is a good raw material for soft drink industry. There have also been reports of jam and jelly production from the pulp following juice extraction (Ohler, 1979). The apple was also found to contain high moisture (80%) and marginal nutrient contents - protein 0.35%; ether extracts 1.8% and total ash 1.9% (Egbe and Sobamiwa, 1989). The cashew nut shell liquid (CNSL), is used widely in brake linings of motor vehicles, paints, varnishes and laminated products (Murthy and Sivasambari, 1985). It is also used as a plywood adhesive (Akaranta *et al.*, 1996) and as a material for increased the tensile properties, as flame retardants of natural rubber (Menon, 1997) and as a long-life, highly bioactive, antifouling coating for marine vessels (Panda and Panda, 1991). CNSL and other extracts from the shell are larvicidal (Carrara *et al.*, 1984; Evans and Raj, 1988) molluscicidal (Casadei *et al.*, 1984; Kubo *et al.*, 1986) and antifungal and antibacterial (Echendu, 1991; Weerasena *et al.*, 1993).

In Nigeria, cashew, like most tree crops hosts a wide range of pests and diseases. Insect pests infest its various parts including roots, stem, branches, flowers, inflorescence and the pseudo-apples (Eguagie, 1972; 1973; 1974), but only a few of these insect species cause economic damage to this crop (Omole, 1972). The occurrence of *Dysdercus superstitionus* (cotton stainer), and *Leptoglossus membranaceus* as new pests of cashew (Olunloyo, 1989), and the recent observations of "sudden death" of mature cashew trees due to infestation by *Plocaederus ferrugineus* L. (Coleoptera: Cerambycidae) (Asogwa *et al.*, 2008), was an indication of a continuous increase in the number and species of insects infesting cashew plantation with time in Nigeria.

<sup>1</sup> This work therefore is an attempt to identify the host plant range and to describe the morphology of the various life stages of an emerging insect pest of cashew, *Plocaederus ferrugineus* in Nigeria.

## MATERIALS AND METHODS

**Survey of cashew plots:** An assessment of cashew germplasm plot (S/GP), which is a 2.5 hectare cashew plot within CRIN headquarters, was carried out to determine the intensity of infestation by the trunk and root borer, *P. ferrugineus*. The layout design of the germplasm experimental plots was a Randomized Complete Block Design with 5 blocks of 0.5 ha each. All other tree/shrub stands (over 23 species) found within the vicinity and adjacent plots were closely observed for any signs of infestation by the borer. The identity of each tree was confirmed at the herbarium of the Forestry Research Institute of Nigeria (FRIN), Ibadan (Table 1).

Table 1. The infestation rate of *Plocaederus ferrugineus* on the various trees/shrubs within the cashew germplasm plots

| Scientific name                   | Family               | Infestation rate (%)*   |
|-----------------------------------|----------------------|-------------------------|
| <i>Anacardium occidentale</i>     | <b>Anacardiaceae</b> | <b>36.3<sup>b</sup></b> |
| <i>Rothmannia longiflora</i>      | Rubiaceae            | 0 <sup>c</sup>          |
| <i>Blighia sapida</i>             | Sapindaceae          | 0 <sup>c</sup>          |
| <i>Alchornea cordifolia</i>       | Euphorbiaceae        | 0 <sup>c</sup>          |
| <i>Combretum sp</i>               | Combretaceae         | 0 <sup>c</sup>          |
| <i>Lecaniodiscus culpanioides</i> | Combretaceae         | 0 <sup>c</sup>          |
| <i>Cola gigantea</i>              | Sterculiaceae        | 0 <sup>c</sup>          |
| <i>Ceiba petandra</i>             | Bombacaceae          | 0 <sup>c</sup>          |
| <i>Albizia zygia</i>              | Mimosoideae          | 0 <sup>c</sup>          |
| <i>Lonchocarpus sericeus</i>      | Papilionoides        | 0 <sup>c</sup>          |
| <i>Dialium guineense</i>          | Caesalpiniaceae      | 0 <sup>c</sup>          |
| <i>Antiaris africana</i>          | Moraceae             | 0 <sup>c</sup>          |
| <i>Gliricidia sepium</i>          | Papilionoides        | 0 <sup>c</sup>          |
| <i>Rauvolfia vomitoria</i>        | Apocynaceae          | 0 <sup>c</sup>          |
| <i>Milicia exelsa</i>             | Moraceae             | 0 <sup>c</sup>          |
| <i>Ficus exasperata</i>           | Moraceae             | 0 <sup>c</sup>          |
| <i>Sterculia tragacantha</i>      | Sterculiaceae        | 0 <sup>c</sup>          |
| <i>Baphia nitida</i>              | Papilionoides        | 0 <sup>c</sup>          |
| <i>Peltophorum pterocarpum</i>    | Caesalpiniodes       | 0 <sup>c</sup>          |
| <i>Pterocarpus osun</i>           | Papilliionoides      | 0 <sup>c</sup>          |
| <i>Cola millenii</i>              | Sterculiaceae        | 0 <sup>c</sup>          |
| <i>Spondias mombin</i>            | <b>Anacardiaceae</b> | <b>89.6<sup>a</sup></b> |
| <i>Newbouldia laevis</i>          | Bignonaceae          | 0 <sup>c</sup>          |
| <i>Psidium guajava</i>            | Myrtaceae            | 0 <sup>c</sup>          |
| <i>Ricinodendron heudelotii</i>   | Euphorbiacea         | 0 <sup>c</sup>          |

Rates with the same superscript are not significantly different ( $P > 0.05$ ) by DMRT.

**Removal of affected cashew trees and the alternate hosts:** The few affected cashew stands were carefully cut down with a saw machine and the stump and root extension carefully exhumed. All the trees and shrubs that exhibited similar infestation symptoms in the 2.5 hectare cashew germplasm plot were cut down and excavated. The total number of the different trunk and the roots excavated were recorded before they were taken to the laboratory for further assessment.

**Morphometrics characteristics of the various stages of the pest:** The excavated cashew and *Spondias mombin* stumps were carefully dissected to expose the various developmental stages of the insects (larvae, pupae and adults), which were collected and assessed. Twenty random samples of each of the developmental stages were selected and used for this study. The adults and larvae were gently killed by placing them in a kilnerjar bottles containing an absorbent cotton wool soaked in ethyl acetate at the bottom. Measurement of morphometric characteristics such as, body length, antennal length, head capsule length, head capsule width, thorax length, thorax width, abdominal length, abdominal width, last abdominal sternum width, elytra length, elytra width, membranous wing length and width was carried out with a pair of caliper.

## RESULTS

Table 1 shows the infestation rate of *Plocaederus ferrugineus* on the various trees/shrubs within the cashew germplasm plots. Observations revealed that fewer stands of cashew were affected by the trunk and root borer resulting in an infestation rate of 36.3%. Out of the 24 species of trees and shrubs found within the germplasm plots, only *Spondias mombin* was found to be riddled with borers. 89.6% of the *S. mombin* assessed came down with different levels of attack by the borer, which was significantly different ( $P < 0.05$ ) from their infestation rate on cashew.

The mean body morphometric measurements of *Plocaederus ferrugineus* larvae were shown in Table 2. The larva (grub) of *P. ferrugineus* has a curled whitish body with wrinkled skin and dark brown head capsule with a mean body length of 2.72mm. The width of the larvae tapers down from the anterior to the posterior end of the body with an anterior width of 0.74mm, which decreased to 0.58mm for both the mid point and posterior width. The larvae however were higher at the midpoint (0.49mm) when compared to the anterior height (0.45mm) and posterior height (0.43).

Table 2. Mean body morphometric measurements of *Plocaederus ferrugineus* larvae (mm)\*

| Parameters              | Mean**             | SD    | SE    | VAR   |
|-------------------------|--------------------|-------|-------|-------|
| Length of body          | 2.724 <sup>a</sup> | 1.189 | 0.266 | 1.414 |
| Height of anterior end  | 0.447 <sup>b</sup> | 0.162 | 0.036 | 0.026 |
| Width of anterior end   | 0.735 <sup>b</sup> | 0.255 | 0.057 | 0.065 |
| Height of posterior end | 0.433 <sup>b</sup> | 0.131 | 0.029 | 0.017 |
| Width of posterior end  | 0.582 <sup>b</sup> | 0.182 | 0.041 | 0.033 |
| Height of mid point     | 0.489 <sup>b</sup> | 0.200 | 0.045 | 0.040 |
| Width of mid point      | 0.582 <sup>b</sup> | 0.207 | 0.046 | 0.043 |

\*Each value represents mean of 20 replicates

\*\*Means with the same superscript are not significantly different ( $P > 0.05$ ) by DMRT.

Table 3. Mean body morphometric measurements of *Plocaederus ferrugineus* pupae (mm)\*

| Parameters              | Mean**             | SD    | SE    | VAR   |
|-------------------------|--------------------|-------|-------|-------|
| Length of body          | 3.660 <sup>a</sup> | 0.469 | 0.105 | 0.220 |
| Height of anterior end  | 1.499 <sup>b</sup> | 0.218 | 0.049 | 0.047 |
| Width of anterior end   | 1.037 <sup>b</sup> | 0.219 | 0.049 | 0.048 |
| Height of posterior end | 1.452 <sup>b</sup> | 0.127 | 0.029 | 0.016 |
| Width of posterior end  | 1.044 <sup>b</sup> | 0.208 | 0.047 | 0.043 |

\*Each value represents mean of 20 replicates

\*\*Means with the same superscript are not significantly different ( $P > 0.05$ ) by DMRT.

The mean body morphometric measurements of *Plocaederus ferrugineus* pupae were shown in Table 3. The pupa is a calcareous cocoon buried in the frass and soft wood tissue of the trunk/roots. An average pupal length of 3.66mm was recorded. However, the mean widths of the pupae at the anterior and posterior ends were not different as both measured 1.04mm. Also the height of the pupae at the anterior end (1.50mm) and posterior end (1.45mm) followed the same pattern as they were not significantly different.

The mean body morphometric measurements of adult *Plocaederus ferrugineus* were shown in Table 4. The adult *P. ferrugineus* is a medium sized dark grey beetle, 2.91cm long and 0.92cm wide at the base of the

abdomen. They have longer antennae (3.58cm), which is significantly longer ( $P < 0.05$ ) than their body (2.91cm). The head, thorax and the elytra are dark brown or almost black. Other morphometric parameters (length of head capsule, width of head capsule and width of elytra; length and width of thorax; width of abdomen, and width of membranous wings) were found to overlap considerably. The lowest value of 0.22cm was recorded for the last abdominal sternum.

Table 4. Mean body morphometric measurements of adult *Plocaederus ferrugineus* (cm)\*

| Parameters                      | Mean**             | SD    | SE    | VAR   |
|---------------------------------|--------------------|-------|-------|-------|
| Length of antennae              | 3.575 <sup>a</sup> | 0.653 | 0.146 | 0.426 |
| Length of head capsule          | 0.590 <sup>g</sup> | 0.112 | 0.025 | 0.013 |
| Width of head capsule           | 0.455 <sup>g</sup> | 0.132 | 0.029 | 0.017 |
| Length of abdomen               | 1.235 <sup>e</sup> | 0.303 | 0.068 | 0.092 |
| Width of abdomen                | 0.830 <sup>f</sup> | 0.169 | 0.378 | 0.289 |
| Length of thorax                | 0.950 <sup>f</sup> | 0.100 | 0.022 | 0.010 |
| Width of thorax                 | 0.915 <sup>f</sup> | 0.127 | 0.028 | 0.016 |
| Length of body                  | 2.905 <sup>b</sup> | 0.369 | 0.083 | 0.136 |
| Length of elytra                | 2.035 <sup>d</sup> | 0.173 | 0.386 | 0.030 |
| Width of elytra                 | 0.515 <sup>g</sup> | 0.081 | 0.018 | 0.007 |
| Length of membranous wing       | 2.540 <sup>c</sup> | 0.302 | 0.067 | 0.091 |
| Width of membranous wing        | 0.915 <sup>f</sup> | 0.150 | 0.033 | 0.022 |
| Width of last abdominal sternum | 0.220 <sup>h</sup> | 0.062 | 0.013 | 0.004 |

\*Each value represents mean of 20 replicates of mixed adult males and females

\*\*Means with the same superscript are not significantly different ( $P > 0.05$ ) by DMRT.

## DISCUSSIONS

*Plocaederus ferrugineus*, which is a serious pest has been reported in India (Abraham, 1958; Pillai *et al.*, 1976; Bhaskara Rao, 1998 ; Mohapatra and Mohapatra, 2004) and Sri Lanka (Gerini, 1976), attacking trunks and roots of the cashew tree. Other related species are *P. consocium*, *P. obesus* and *Bactocera rufomaculata* (Abraham, 1958; Pillai *et al.*, 1976). When caught, the beetle makes an indignant squeaking sound by rubbing the prothorax and mesothorax together.

The establishment of cashew in South Western Nigeria does not often require clean weeding, but line weeding or cutting of traces at a recommended spacing between the lines. The belt of vegetation left is to provide the necessary microclimate for the transplants of cashew, especially when rain ceases. *S. mombin* was found not to be just an “alternate host” of the pest but could be referred to as the “preferred host plant”, as over eighty nine percent (89.6%) of the assessed plants came down with different levels of attack by the borer, which was significantly different ( $P < 0.05$ ) from their infestation rate on cashew. All the other plant species (trees and shrubs) within the vicinity of the germplasm plots were not infested with the borers.

The identification and knowledge of the location of the alternate hosts should facilitate the formulation of effective control strategies for the pest. For example, spot sprays of the insect on the alternate host plants may reduce its population in the environment. Also cutting down of such host plants within the vicinity will reduce or eliminate sources of reinfestation of cashew trees. Various authors have reported the presence and activities of other related borers in the same Order on tree crops. For example, Daramola (1978) and Ojo (1981) reported that the larvae of *Tragocephala castmia* Thoms. (Coleoptera: Cerambycidae) were found to bore into the stems of kola and cocoa trees. Ndubuaku (1989) reported *Phosphorus virescens* and *Phosphorus gabonator* (Coleoptera: Cerambycidae) as two species of stem borers affecting kola trees, which have been occasionally seen in cocoa farms.

Ecological studies often require reliable methods of segregating the sexes of the insect pest under investigation in the field. The morphological characters were usually assessed in some cases to separate the sexes. However, external physical observation of the last abdominal segment of beetles in the same order, like *Analeptes trifasciata* was the simplest and reliable means of segregating the sexes. The females were usually found to possess a mid ventral structure on their last abdominal sternum, which was absent in the males. However, this

structural contrast needs a confirmatory test, which involves dissecting the adults and is therefore being used as a satisfactory criterion for segregating the sexes. This is yet to be done in the case of *P. ferrugineus* a new emerging insect pest of cashew in Nigeria. A mixed adult of both sexes was therefore used to carry out this preliminary study.

## REFERENCES

- Abraham, E. V. 1958. Pests of cashew (*Anacardium occidentale*). *Indian Journal of Agric Sciences (India)*. 28(4): 531-543
- Akaranta, O.; W. Donbebe and T.O. Odozi. 1996. *Plywood adhesives based on red-onion-skin extract modified with cashew nut-shell liquid*. *Bioresource Technol.* 56: 279-280
- Asogwa, E.U.; J.C. Anikwe; T.C.N. Ndubuaku and F.A. Okelana. 2008. *Distribution and damage characteristics of an emerging insect pest of cashew, Plocaederus ferrugineus L. (Coleoptera: Cerambycidae) in Nigeria: a preliminary report*. *African Journal of Biotechnology*, Vol. 7(24) Accepted Dec. 17, 2008 (In press).
- Bhaskara Rao, E.V.V. 1998. *Integrated production practices of cashew in India*. *RAP FAO Corporate Document Repository*. Available online at [www.fao.org/docrep/005/ac451e/ac451e04.htm](http://www.fao.org/docrep/005/ac451e/ac451e04.htm). 12pp.
- Carrara, G.; G.C. Munoz and L. Damho. 1984. *Larvicidal effect of cashew nut husk bagasse: Its possible use in malariology in the antivector control programmes*. *Revista Medica de Mocambique*. 2: 78-82
- Casadei, E.; S. Bruheim and T. Latis. 1984. *Active substances in cashew nut shell with molluscicidal activity: Possible use in Schistosomiasis control programmes*. *Revista Medica de Mocambique*. 2: 35-39
- Deckers, J.; E. Cundall; S.H. Shomari; A. Ngatunga and G. Bassi. 2001. *Nut crops. Cashew*. In: *Romain H. Raemaekers (Editor), Crop Production in Tropical Africa*. Pp. 691-700. Goekint Graphics nv. Belgium.
- Daramola A.M. 1978. Insect pests of kola in Nigeria. *Res. Bull.* No.3: CRIN, Ibadan. 33 pp
- Echendu, T.N.C. 1991. *Ginger, Cashew and Neem as surface protectants of cowpeas against infestation and damage by Callosobruchus maculatus*. *Tropical Science*. 31: 209-211
- Egbe, N.E. and O. Sobamiwa. 1989. *Utilization of Cocoa, Kola, Coffee, Cashew and Tea In Nigeria*. In *Progress in Tree Crop Research*. CRIN, Ibadan - Nigeria. Pp. 217-224
- Eguagie, W.E. 1972. *Insects associated with cashew Anacardium occidentale in Nigeria*. *Ann, Rep. CRIN, Ibadan, Nigeria*. Pp.134-137
- Eguagie, W.E. 1973. *Insect associated with cashew - A. occidentale in Nigeria*. *CRIN Ann. Rep.* Pp. 270-273
- Eguagie, W.E. 1974. *Insects associated with cashew-A. Occidentale in Western Nigeria*. *CRIN. Ann. Rep.* Pp. 128-130.
- Evans, D.A. and R.K. Raj. 1988. *Extracts of Indian plants as mosquito larvicides*. *Indian J. Med. Res.* 88: 38-41
- Gerini, V. 1976. *I coleotteri presenti sull'anacardio con particolare riferim ento a P. ferrugineus nello Sri-Lanka*. *Rivista di Agricoltura Subtropicale e Tropicale (Italy)*. 70(7-9/10-12): 185-190
- Hammed, L.A.; A.R. Adedeji; E.U. Asogwa and O.S. Ibiremo. 2007. *Constraints to cashew production in Nigeria. A paper presented at the cashew stake-holders meeting organised by the African Cashew Alliance (ACA) held at IITA, Ibadan, Nigeria*. January 26, 2007. 12pp
- Kubo, I.; S. Komatsu and M. Ochi. 1986. *Molluscicides from cashew, Anacardium occidentale and their large scale isolation*. *J. Agric Food Chem.* 41: 1012-1015.
- Menon, A.R.R. 1997. *Flame-retardant characteristics of natural rubber modified with a bromo derivative of phosphorylated cashew nut shell liquid*. *J. fire Science.* 15: 3-13
- Mohapatra, L.N. and R.N. Mohapatra. 2004. *Distribution, intensity and damage of cashew stem and root borer P. ferrugineaus in Orissa*. *Indian Journal of Entomology*. (Vol 66) (No. 1): 4-7
- Murthy, B.G.K. and M.A. Sivasamban. 1985. *Recent trends in CNSL utilization. Cashew Research and development: Proceedings of the International Cashew Symposium, Cochin, Kerala, India. March 12-15, 1979*. Pp. 201-207

- Ndubuaku, T.C.N. (1989). *Economic insect pests of Kola. In: Progress in Tree Crops Research. 2<sup>nd</sup>. CRIN, Ibadan, Nigeria. Pp. 115-126*
- Ohler, J.G. 1967. *Cashew Growing. Tropical Abstracts (The Netherlands). 22(1): 1-9*
- Ohler, J.G. 1979. *Cashew. Koninklijk Instituut Voor de Tropen Amsterdam. 260pp*
- Ojo. A. A 1981. *Insect pests and cocoa production in Nigeria. Proceeding of the Nigeria Cocoa Board symposium held at the University of Ibadan, Nigeria. August 20, 1980. Pp. 42-49*
- Olunloyo, O. A. 1989. *Important diseases of cashew and their control. In Progress in Tree Crops Research. CRIN, Ibadan, Nigeria. Pp. 175-190*
- Omole, M.M. 1972. *Insects associated with cashew A. occidentale in Nigeria. Annual Report, CRIN. Pp 134-137*
- Opeke, L.K. 1987. *Tropical Tree Crops. Spectrum Books Ltd, Ibadan. 326pp*
- Panda, R. and H. Panda. 1991. *Antifouling coatings based on cashew nut shell liquid (CNSL) modified resin. Paint Ink Int. 4: 30-32*
- Pillai G.B.; O.P. Dubey and V. Singh (1976). *Pests of cashew and their control in India: A review of current status. J. Plant Crops 4: 37-50.*
- Togun, A. 1977. *A review of the prospect of cashew industry. 39pp*
- Venkataramah, T.M. 1976. *Cashew nut production and processing – Nigeria agronomic aspect of cashew nut production. Unpublished paper submitted to CRIN. 39pp*
- Weerasena, O.V.D.S.J.; A.S. Amarasekara and R.L.C. Wijesundera. 1993. *Fungicidal activity of synthetically modified cashew nut shell. J. Nat Sc. Coun. Sri-Lanka. 21: 253-258*