

EFFECT OF *Trichoderma* ON SEED GERMINATION AND SEEDLING PARAMETERS IN CHILIM.S. ISLAM¹, M.A. RAHMAN², S.H. BULBUL³, AND M.F. ALAM⁴¹PSO, Hybrid Rice Project, BRRI, Gazipur-1701; ²Department of Botany, University of Rajshahi, Rajshahi-6205, Bangladesh; ³Researcher, "BAS-BRRI-34"- Project, Plant Breeding Division, Bangladesh Rice Research Institute, Gazipur-1701; ⁴Biotechnology and Microbiology laboratory, Department of Botany, University of Rajshahi, Rajshahi-6205, Bangladesh.

Corresponding author & address: Dr. Md. Shafiqul Islam, PSO, Hybrid Rice Project, BRRI, Gazipur-1701, shafiqpabna@yahoo.com

Accepted for publication on 15 December 2010

ABSTRACTIslam MS, Rahman MA, Bulbul SH, Alam MF (2011) Effect of *Trichoderma* on seed germination and seedling parameters in chili. *Int. J. Expt. Agric.* 2(1), 21-26.

T. virens IMI-392430, *T. pseudokoningii* IMI-392431, *T. harzianum* IMI-392432, *T. harzianum* IMI-392433 and *T. harzianum* IMI-392434 were evaluated for their potentiality on seed germination and seedling parameters in chili both laboratory and field conditions. Chili seeds were coated with spore suspension of each test strains of *Trichoderma* supplemented with 2 % of starch (w/v) as an adhesive. Seed germination percentages and the vigour index were significantly ($P \leq 0.05$) affected by the application of different strains of *Trichoderma*. Among the five *Trichoderma* strains, *T. harzianum* IMI-3924332 gave the highest germination percentage followed by *T. harzianum* IMI-3924333, *T. harzianum* IMI-3924334, *T. virens* IMI-392430 and *T. pseudokoningii* IMI-392431 treatment both in laboratory and field conditions, respectively while control decrease these value. Chili seeds also gave the highest vigour index values with *T. harzianum* IMI-3924332 which confirmed to better germination. Seed treatment with *T. harzianum* IMI-3924332 can be useful to enhance the germination of chili seeds as well as reduce to delayed germination. Further investigations however are required to study *in vivo* effect of *Trichoderma* strains on morphological and physiological characteristics in chili plant and fruit production.

Key words: *Trichoderma*, spore suspension, germination percentages, vigour index, chili**INTRODUCTION**

Chili (*Capsicum annum* L.) is one of the most important spice crops in the world and grown in all seasons and areas of Bangladesh. The average yield of chili is 0.042 t ha⁻¹ which is very low as compared to the yield of other chili growing countries of the world (Anonymous 2003). Delayed and erratic germination of chili seeds is one of the reasons of low yield of chili. There are many factors responsible for the delayed and erratic germination of chili seeds. Among the various factors diseases are predominant. Fungal diseases play a vital role in reducing the germination of chili. Water imbibitions are first step in the seed germination. But crop field may lack adequate moisture content for the same, so poor and delayed germination occurs. This leads to suffocation, resulting in delayed and poor germination as well as weak seedling growth (Heydecker 1977).

Some authors have defined *Trichoderma* strains as plant symbiont opportunistic avirulent organisms, able to colonize plant roots by mechanisms similar to those of mycorrhizal fungi and to produce compounds that stimulate growth and plant defense mechanisms. Crop productivity in fields can increase up to 30% after the addition of *Trichoderma hamatum* or *Trichoderma koningii* (Benítez *et al.* 2004). *Trichoderma* strains that produce cytokinin-like molecules, e.g. zeatyn and gibberellin GA3 or GA3-related have been recently detected. Moreover, recent studies have indicated that these fungi also induce localized or systemic resistance systems in plants (Yedidia *et al.* 1999; Howell 2003). Thus, the variety of effects indicates that these beneficial fungi have multiple modes of action. Therefore, this study was design to find out the effect of different *Trichoderma* strains on the germination percentage rate and seedling parameters of chili seeds both in laboratory and field conditions.

MATERIALS AND METHODS**Sources of *Trichoderma* strains**

Five *Trichoderma* strains viz. *T. virens* IMI-392430, *T. pseudokoningii* IMI-392431 and *T. harzianum* IMI-392432, *T. harzianum* IMI-392433 and *T. harzianum* IMI-392434 were used in this study which was collected from Biotechnology and Microbiology Laboratory, Department of Botany, Rajshahi University, Bangladesh. These strains were isolated and identified from decomposed garbage and soil by Rahman (2009) and were verified by CABI Bioscience, Surrey, U.K.

Spore suspension preparation of *Trichoderma* strains

Mycelial disc (1.2 cm diam.) of *Trichoderma* strains were obtained from 4-5 days-old culture and transferred to 50 ml PDA in a 250-ml conical flask separately incubated at 28 °C for 4-5 days. At the end of the incubation period 30 ml of sterile distilled water was added to each culture flask and the flasks were shaken at 50 rpm for 30 min in an orbital shaker. Then the content of each conical flask was filtrate through sterile muslin cloth. The filtrate with the spores was collected and a concentration of spore suspension was adjusted to 5×10^5 conidia/ml by use of a hemacytometer under a light microscope.

Seed selection and treatment

The seed of chili Variety Bogra Local was used and collected from Spice Research Centre, Bogra, Bangladesh. The chili seeds were one year old and had been stored at 5°C. Standard germination of the seeds was 98%. Seeds with no cracks or

other visible deformations were selected and surface sterilized for 10 minutes with 1% sodium hypochlorite solution. Seeds were then rinsed three times with sterile distilled water and air dried. A seed coating was prepared from spore suspension supplemented with 2% of starch (w/v) as an adhesive. Dry chili seeds were dipped in seed coating suspension (5×10^5 spores/ml) for each *Trichoderma* strains for 1-2 minutes. For untreated control seeds were dipped in 2% starch suspension and for water control seeds were dipped in sterilized distilled water. Seeds were air dried inside the laminar air flow hood. For *in vitro* experiment, treated seeds were placed in Petri plate's lines with two layers of Whatman filter paper soaked in sterilized distilled water and incubated at 25°C under dark condition and for field experiment, treated seeds were sown separately in pot soils where the soil was previously inoculated with respective *Trichoderma* strains (5×10^5 spore/ml) and for control treatment, treated seeds (treated with 2% starch and water) were sowing un inoculated soil in pot. At least ten seeds were sown in each pot. Seed germination percentages and vigour index was recorded after 3 to 8 days. Vigour index for each treatment was determined using the following formula developed by Abdul-Baki and Anderson (1973).

Vigour index = [Mean of root length (cm) + Mean of shoot length (cm)] \times percentages of seed germination.

Collection and preparation of soil for field experiment

For field experiment soil was collected from the Botanical garden of Rajshahi University Campus, Bangladesh and sterilized with Formaldehyde (Formalin: Water; 1:5 V/v). After 30 days of sterilization, soils were put in the earth pot of 12 inches height and 8 inches wide. For minimize losses of excess water 2 cm hole was made from the bottom of the pot.

Experimental design and data analysis

All experiments were established as a randomized block design with four replicates and ten chili seeds were used in each replicates. Data on germination percentages and vigour index were recorded after 3 to 8 days and statistically analyzed with the help of computer package program SPSS (SPSS Inc., Chicago, IL, USA) and also tested by DMRT.

RESULTS AND DISCUSSION

The effect of five *Trichoderma* strains on seed germination and seedling parameters of chili both in laboratory and field conditions the results are presented in Fig 1-5. Statistical analysis of figure showed significant differences in treatments at $P \leq 0.05$ levels. Results showed that all *Trichoderma* strains were found effective to enhance the germination percentage compared to control. However among the five *Trichoderma* strains, *T. harzianum* IMI 392432 exhibited significantly enhancement of germination percentage in chili seeds both in laboratory and field conditions followed by *T. harzianum* IMI 392433, *T. harzianum* IMI 392434, *T. virens* IMI 392430 and *T. pseudokoningii* IMI 392431 (Fig 1 and 5), while control (treated with 2% starch and water) significantly decreased these values. This strains also showed earliest highest seed germination (100%) at five and six days compared to the control both laboratory and field conditions, respectively. In controls (treated with 2% starch and water), both laboratory and field conditions showed worst germination percentage and vigour index. Some landmarks along the way include the discoveries that these fungi frequently increase plant growth and productivity (Harman 2006; Manju and Mall, 2008). In this study, five *Trichoderma* strains gave early germination as well as highest germination percentage which have also been reported by many workers in different plants (Hanson 2000; Mishra and Sinha, 2000; Oyarbide *et al.* 2001) and numerous other species such as *T. longipile* and *T. tomentosum* have been shown to promote plant growth (Rabeendran *et al.* 2000). Studies have been confirmed in case of *T. harzianum* and *T. viridi* to enhanced seed germination root and shoot length (Dubey *et al.* 2007) as well as increasing the frequency of healthy plants, and boosting yield (Rojoa *et al.* 2007). In a similar study Chaur-Tsuen Lo and Chien-Yih Lin (2002) screened *Trichoderma* strains on plant growth and root growth of bitter melon, loofah and cucumber and noted that *Trichoderma* strains significantly increased of 26 to 61% in seedling height, 85-209% in root exploration, 27-38% in leaf area and 38 to 62% in root dry weight after 15 days of sowing. Methanol extract of *T. harzianum* and *T. viridi* significantly improved various growth parameters of okra (Prasad and Anes, 2008). Vigour index (VI) was also significantly affected by the application of different *Trichoderma* strains both in laboratory and field conditions (Fig 4). The results related to vigour index showed similar differences as in germination percentages. Seed treatment with *Trichoderma* strains increased vigour index compared to control. The highest VI values were recorded both in laboratory and field conditions when the chili seed were treated with *T. harzianum* IMI 392432. The lowest vigour index was recorded in control. Mukhtar (2008) investigated that seed treatment with *T. harzianum* gave the highest germination index in okra and *T. harzianum* can be useful to enhance the germination percentage as well as reduce lose due to delayed germination of okra seeds. Begum *et al.* (2010) were evaluated five *Trichoderma* strains to assay their efficacy in suppressing *Alternaria* fruit rot disease of chili and promoting chili plant growth and yield and observed that application of *T. harzianum* IMI 392432 significantly suppressed the disease and improved highest seed germination percentage, vigour index, growth and yield.

CONCLUSION

The present study concludes that *Trichoderma* species have potential to enhance the germination in chili seeds which can be useful to enhance the germination percentage of chili seeds besides reducing losses due to delayed germination. Further

investigations are required to study *in vivo*, effects of these fungi on the morphological and physiological characteristics in chili plant and fruit production.

REFERENCES

- Abdul-Baki A, Anderson JD (1973) Vigour determination of Soyabean seed by multiple criteria. *Crop Sci* 13, 630-633.
- Anonymous (2003) Monthly Statistical Bulletin, Bangladesh. Bangladesh Bureau of Statistics, August 2003. p 54.
- Begum MF, Rahman MA, Alam MF (2010) Biological control of *Alternaria* fruit rot of chili by *Trichoderma* species under field conditions. *Mycobiology* 38(2), 113-117.
- Benítez T, Rincón AMM, Limón C, Codón AC (2004) Biocontrol mechanisms of *Trichoderma* strains. *International Microbiology* 7, 249-260.
- Chaur-Tsuen L, Chien-Yih L (2002) Screening strains of *Trichoderma* spp. for plant growth enhancement in Taiwan. *Plant Pathology Bulletin* 11, 215-220.
- Dubey SC, Suresha M, Singha B (2007) Evaluation of *Trichoderma* species against *Fusarium oxysporum* f. sp. ciceris for integrated management of chickpea wilt. *Biological Control* 40, 118-127.
- Hanson LD (2000) Reduction of *Verticillium* wilt symptoms in cotton following seed treatment with *Trichoderma virens*. *J. Cotton. Sci.* 4:224-231.
- Harman GE (2006) Overview of mechanisms and uses of *Trichoderma* spp. *Phytopathology* 96, 190-194.
- Heydecker W (1977) Stress and seed germination: and agronomic view. In the physiology and biochemistry of seed dormancy and germination, A. A Khan, ed. Amsterdam: North Holland Publishing Company.
- Howell CR (2003) Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: The history and evolution of current concepts. *Plant Dis.*, 87, 4-10.
- Manju S, Mall TP (2008) Efficacy of *Trichoderma* species on *Phytophthora dresceleri* f. sp. *cajani* of Pigeon pea. *Ann. Plant Prot. Sci.* 16, 162-164.
- Mishra DS, Sinha AP (2000) Plant growth promoting activity of some fungal and bacteria agents on rice seed germination and seedling growth. *Tropical Agric.* 77, 188-191.
- Mukhtar I (2008) Influence of *Trichoderma* species on seed germination in okra. *Mycopath* 6(1&2), 47-50
- Oyarbide F, Osterrieth ML, Cabello M (2001) *Trichoderma koningii* as a biomineralizing fungous agent of calcium oxalate crystals in typical Argiudolls of the Los Padres Lake natural reserve (Buenos Aires, Argentina). *Microbiol. Res.* 156, 113-119.
- Prasad D, Anes KM (2008) Effect of metabolites of *Trichoderma harzianum* and *T. viride* on plant growth and meloidogyne incognita on okra. *Ann. Plant Prot. Sci.* 16, 461-465.
- Rabeendran N, Moot DJ, Jones EE, Stewart A (2000) Inconsistent growth promotion of cabbage and lettuce from *Trichoderma* isolates. *New Zealand Plant Prot.* 53, 143-146.
- Rahman MA (2009) Screening of *Trichoderma* spp. and their efficacy as a bio conversion agent of municipal solid waste through appropriate technique of solid state fermentation. PhD Thesis. Department of Botany, University of Rajshahi, Rajshahi-6205, Bangladesh.
- Rojoa FG, Reynosa MM, Fereza M, Chulze SN, Torres AM (2007) Biological control by *Trichoderma* species of *Fusarium solani* causing peanut brown root rot under field conditions. *Crop Prot.* 26, 549-555.
- Yedia I, Benhamou N, Chet I (1999) Induction of defense responses in cucumber plants (*Cucumis sativus* L.) by the biocontrol agent *Trichoderma harzianum*. *Appl. Environ. Microbiol.*, 65, 1061-1070.




Fig 1. Effect of seed treatment with *Trichoderma* strains on the germination percentage of chili seeds in *in vitro* condition. Vertical bars show standard error of means of four replicates. Bar marked by the same letters are not significantly different ($P < 0.05$) by DMRT analysis




Fig 2. Effect of seed treatment with *Trichoderma* strains on shoot length of chili seeds in *in vitro* condition. Vertical bars show standard error of means of four replicates. Bar marked by the same letters are not significantly different ($P < 0.05$) by DMRT analysis.




Fig 3. Effect of seed treatment with *Trichoderma* strains on root length of chili seeds in *in vitro* condition. Vertical bars show standard error of means of four replicates. Bar marked by the same letters are not significantly different ($P < 0.05$) by DMRT analysis.




Fig 4. Effect of seed treatment with *Trichoderma* strains on vigour index of chili seeds in *in vitro* condition. Vertical bars show standard error of means of four replicates. Bar marked by the same letters are not significantly different ($P < 0.05$) by DMRT analysis.




Fig 5. Effect of seed treatment with *Trichoderma* strains on the germination percentage of chili seeds in field condition. Vertical bars show standard error of means of four replicates. Bar marked by the same letters are not significantly different ($P < 0.05$) by DMRT analysis.